arXiv:1702.08012v1 [math.RT] 26 Feb 2017

CUSPIDAL CHARACTERS AND AUTOMORPHISMS

GUNTER MALLE

Abstract. We investigate the action of outer automorphisms of finite groups of Lie type on their irreducible characters. We obtain a definite result for cuspidal characters. As an application we verify the inductive McKay condition for some further infinite families of simple groups at certain primes.

1. Introduction

An important open problem in the ordinary representation theory of finite groups of Lie type is to determine the action of outer automorphisms on the set of their irreducible characters, and more generally to determine the irreducible character degrees of the corresponding almost (quasi-)simple groups. While the action of diagonal automorphisms and the corresponding extension problems are well understood by the work of Lusztig, based on the fact that such extensions can be studied in the framework of finite reductive groups, much less is known in the case of field and also of graph automorphisms.

The most elusive situation seems to be the one where irreducible characters not stable under diagonal automorphisms are concerned. In [12] we obtained a certain reduction of this problem to the case of cuspidal characters. This is the situation we solve here by applying methods and results from block theory and Deligne–Lusztig theory (see Section 4):

Theorem 1. *Let G be a quasi-simple group of Lie type. For any cuspidal character ρ of G there is a semisimple character χ in the rational Lusztig series of ρ having the same stabiliser as ρ in the automorphism group of G.*

Observe that the action on semisimple characters is well-understood by the theory of Gelfand–Graev characters, see [15]. For linear and unitary groups this result was obtained by Cabanes and Späth [5] and we use it in our proof, for symplectic groups it follows from recent work of Cabanes–Späth [6] and of Taylor [16]; for the other types it is new. For the proof we first consider quasi-isolated series, see Sections 2 and 3. Here, we connect ρ to χ either via a sequence of Brauer trees, in which case we also obtain information on maximal extendibility (see Corollary 3.3), or of Deligne–Lusztig characters.

As an application we verify the inductive McKay condition for some series of simple groups of Lie type and suitable primes ℓ (see Section 5):

Theorem 2. *Let q be a prime power and $S = {}^2E_6(q)$, $E_7(q)$, $B_n(q)$ or $C_n(q)$ be simple. Let $\ell \equiv 3 \pmod 4$ be a prime with $\ell|(q^2-1)$. Then S satisfies the inductive McKay condition at ℓ. In particular, the inductive McKay condition holds for S at $\ell = 3$.*

Date: February 28, 2017.

2010 *Mathematics Subject Classification.* Primary 20C15, 20C33; Secondary 20G40.

Key words and phrases. McKay conjecture, automorphisms, action on cuspidal characters.

The author acknowledges financial support by ERC Advanced Grant 291512 and by DFG TRR 195.

This is a contribution to a programme to prove McKay's 1973 conjecture on characters of ℓ'-degree based on its reduction to properties of quasi-simple groups, an approach which has recently led to the completion of the proof in the case when $\ell = 2$ (see [12]).

Acknowledgement: I thank David Craven for his remark that the consideration of Brauer trees might be useful, and the Institut Mittag-Leffler, where this remark was made, for its hospitality. I also thank Frank Lübeck for help in compiling Table 2, and Britta Späth and Jay Taylor for helpful comments on an earlier version.

2. Cuspidal characters in classical groups

Throughout the paper we fix the following notation. We let $\mathbf{G}$ be a simple simply connected linear algebraic group over an algebraically closed field of characteristic p with a Frobenius map $F : \mathbf{G} \to \mathbf{G}$ inducing an $\mathbb{F}_q$-structure, and we set $G := \mathbf{G}^F$. Let $\mathbf{G} \hookrightarrow \widetilde{\mathbf{G}}$ be a regular embedding; thus $\widetilde{\mathbf{G}}$ is a group with connected centre and derived subgroup equal to $\mathbf{G}$. For an extension $F : \widetilde{\mathbf{G}} \to \widetilde{\mathbf{G}}$ of the Frobenius endomorphism on $\mathbf{G}$ we let $\tilde{G} := \widetilde{\mathbf{G}}^F$. We choose a group $\widetilde{\mathbf{G}}^*$ dual to $\widetilde{\mathbf{G}}$, with corresponding Frobenius map again denoted by F, and an epimorphism $\pi : \widetilde{\mathbf{G}}^* \to \mathbf{G}^*$ dual to the regular embedding $\mathbf{G} \hookrightarrow \widetilde{\mathbf{G}}$, and we write $\tilde{G}^* = \widetilde{\mathbf{G}}^{*F}$ and $G^* = \mathbf{G}^{*F}$ for the F-fixed points.

2.1. **Cuspidal unipotent characters on Brauer trees.** One crucial tool in our determination of the action of automorphisms is the observation that cuspidal unipotent characters of classical groups lie in blocks of cyclic defect for suitable primes. According to Lusztig's classification, recalled in Table 1, a classical group G has at most one cuspidal unipotent character, which is thus in particular fixed by all automorphisms of G.

G	A_{n-1}	${}^2A_{n-1}$	B_n, C_n	D_n	${}^2D_n$
n	1	$a(a+1)/2$	$a(a+1)$	a^2 even	a^2 odd
label	$()$	$(a, a-1, \ldots, 1)$	$\binom{0\ldots 2a}{-}$	$\binom{0\ldots 2a-1}{-}$	$\binom{0\ldots 2a-1}{-}$
d_G	$-$	$2(2a-1)$	$4a$	$2(2a-1)$	$2(2a-1)$

Table 1. Cuspidal unipotent characters in classical groups

From the hook formula (see e.g. [14]) it follows that the cuspidal unipotent character of G is of defect zero for every Zsigmondy prime of odd degree, as well as for those of even degree $d > d_G$. Recall that for every prime power q and any integer $d > 2$ with $(q, d) \neq (2, 6)$ there exists a *Zsigmondy primitive prime of degree* d dividing $q^d - 1$ but no $q^f - 1$ for $f < d$. We write Φ_d for the dth cyclotomic polynomial over $\mathbb{Q}$.

Lemma 2.1. *Let G be simple of classical type with a cuspidal unipotent character ρ. Then there exist sequences of unipotent characters $\rho = \rho_1, \ldots, \rho_m = 1_G$ of G and of Zsigmondy primes $\ell_i \neq p$ of either odd degree $d > 2$ or even degree $d \geq d_G$ such that ρ_i, ρ_{i+1} lie in the same ℓ_i-block of cyclic defect of G, for $i = 1, \ldots, m-1$, except when $G = D_4(2)$.*

Proof. The claim is clear for type A_0 as here the cuspidal character is the trivial character. For G of type ${}^2A_{n-1}$ there exists a cuspidal unipotent character ρ if and only if $n = a(a+1)/2$ for some $a \geq 2$ (see Table 1). This is labelled by the triangular partition

$\delta_a = (a, \dots, 1)$ of n, which has a unique hook of length $2a-1$. By the hook formula (see e.g. [14]) this implies that $|G|/\rho(1)$ is divisible by $\Phi_d(q)$ exactly once, where $d = 2(2a-1)$. So by [9] ρ lies in an ℓ_1-block of cyclic defect for any Zsigmondy prime divisor ℓ_1 of $q^{2a-1}+1$. (Such a prime ℓ_1 exists unless $(q,a) = (2,2)$, in which case $n = 3$, but ${}^2A_2(2)$ is solvable.) Moreover, the partition $(3a-3, a-3, \dots, 1)$ has the same $2a-1$-core as δ_a, so the unipotent character ρ_2 labelled by it lies in the same ℓ_1-block as $\rho_1 = \rho$. The latter partition has a unique $4a-4$-hook, and arguing as before, we conclude that ρ_2 lies in the same ℓ_2-block of cyclic defect as the character ρ_3 labelled by $(5a-10, a-5, \dots, 1)$, for ℓ_2 a Zsigmondy prime divisor of $q^{4a-4}+1$. Continuing inductively we arrive at $\rho_{2a+1} = 1_G$.

Groups of type B_n and C_n have a cuspidal unipotent character ρ if and only if $n = a(a+1)$ for some $a \geq 1$. This is labelled by the symbol $\binom{0\ \dots\ 2a}{\ \ -\ \ }$, which has a single $2a$-cohook. So ρ lies in a block of cyclic defect for any Zsigmondy prime divisor ℓ_1 of $q^{2a}+1$ (see again [9]). Furthermore this block also contains the unipotent character ρ_2 labelled by $\binom{0\ \dots\ 2a-2}{0\ \ 4a-1}$. This has a unique $4a-2$-hook, whose removal gives $\binom{0\ \dots\ 2a-2}{0\ \ \ 1}$, and adding a suitable $4a-2$-hook we find $\binom{0\ \dots\ 2a-3\ \ 6a-4}{\ \ \ \ \ 0\ \ 1}$, the symbol of a unipotent character ρ_3 lying on the same Brauer tree as ρ_2 for primes ℓ_3 dividing $q^{4a-2}-1$. The Zsigmondy exception $a = 2$ can be avoided by working with 5-hooks instead of 6-hooks in this step. Continuing this way, after $m = a$ steps we arrive at the symbol of $\rho_{a+1} = 1_G$.

For groups of type D_n or ${}^2D_n$, $n \geq 4$, a cuspidal unipotent character exists only if $n = a^2$ for some $a \geq 2$. It is labelled by the symbol $\binom{0\ \dots\ 2a-1}{\ \ -\ \ }$. This has a single $2a-1$-cohook and thus lies in an ℓ_1-block of cyclic defect for ℓ_1 any primitive prime divisor of $q^{2a-1}+1$. (Such a prime ℓ_1 exists unless $(q,a) = (2,2)$, which leads to the stated exception.) This block also contains the character ρ_2 labelled by $\binom{0\ \dots\ 2a-3}{0\ \ 4a-3}$. The latter symbol has a unique $4a-4$-hook; removing this and adding a different one leads to the symbol $\binom{0\ \dots\ 2a-4\ \ 6a-7}{\ \ \ \ \ 0\ \ 1}$ of a unipotent character ρ_3. Again, induction completes the proof. □

2.2. **Constituents of Lusztig induction.** We will also need a result on Lusztig induction from certain 2-split Levi subgroups.

Lemma 2.2. *Let G be quasi-simple of classical type B_n, C_n, D_n or ${}^2D_n$ and ρ be a cuspidal unipotent character of G. Let $\mathbf{L} \leq \mathbf{G}$ be a 2-split Levi subgroup of $\mathbf{G}$*

1. *of twisted type ${}^2A_{n-1}(q).\Phi_2$ in type B_n, C_n; or*
2. *of twisted type ${}^2A_{n-2}(q).\Phi_2^2$ in type D_n or ${}^2D_n$.*

There exist sequences of unipotent characters $\rho_1 = \rho, \dots, \rho_m = 1_G$ of G and $\psi_1, \dots, \psi_{m-1}$ of L such that ρ_i, ρ_{i+1} both occur with multiplicity ± 1 in $R_{\mathbf{L}}^{\mathbf{G}}(\psi_i)$ for $i = 1, \dots, m-1$.

Proof. First consider types B_n and C_n. So $n = a(a+1)$ for some $a \geq 1$, and ρ is parametrised by the symbol $\mathcal{S} = \binom{0\ \dots\ 2a}{\ \ -\ \ }$. Since this symbol has exactly n 1-cohooks, its 1-cocore is trivial and so ρ lies in the principal 2-series. Observe that all unipotent characters of groups of type A are uniform, so that $R_{\mathbf{L}}^{\mathbf{G}}$ can be expressed in terms of Deligne–Lusztig characters $R_{\mathbf{T}}^{\mathbf{G}}$ for various F-stable maximal tori $\mathbf{T}$ of $\mathbf{G}$. The decomposition of Deligne–Lusztig characters has been determined explicitly by Lusztig, and from this our claim can be checked by direct computation.

To do this, we appeal to [2, Thm. 3.2], which shows that Lusztig induction of unipotent characters from 2-split Levi subgroups is, up to signs, the same as induction in the

corresponding relative Weyl groups. The relative Weyl group for the principal 2-Harish-Chandra series of G is $W{=}W(B_n)$, that of L is its maximal parabolic subgroup $W_L = \mathfrak{S}_n$. Determination of the 2-quotient of the symbol $\mathcal{S}$ shows that it corresponds to the character with label $(\delta_a;\delta_a)$ of W, with $\delta_a = (a,\dots,1)$ the triangular partition. Let ψ be the unipotent character of L parametrised by the partition $2\delta_a = (2a, 2a-2, \dots, 2)$. The constituents of $\mathrm{Ind}_{W_L}^W(2\delta_a)$ are exactly those bipartitions whose parts (including zeroes) can be added up so as to obtain the partition $2\delta_a$, with multiplicity one if this is possible in a unique way. Thus it contains $(\delta_a;\delta_a)$ exactly once, but also $(2a, 2a-4, \dots; 2a-2, 2a-6, \dots)$. This in turn is contained once in $\mathrm{Ind}_{W_L}^W((4a-2, 4a-10, \dots))$, and so on. Continuing in this way we reach $(a(a+1);-)$ which parametrises the trivial character.

For G of type D_n, a cuspidal unipotent character exists if $n = a^2$ for some even $a \geq 2$, and it is labelled by the symbol $\binom{0\cdots 2a-1}{-}$. Again, this character lies in the principal 2-series of G, and by [2, Thm. 3.2] the decomposition of $R_{\mathbf{L}}^{\mathbf{G}}$ can be computed in the relative Weyl groups $W_L = \mathfrak{S}_{n-1} \leq W = W(D_n)$. Here, ρ corresponds to the character of W with label $(\delta_a;\delta_{a-1})$. Let $W_1 = \mathfrak{S}_n \leq W(D_n)$ be a maximal parabolic subgroup of $W(D_n)$ containing W_L. For ψ labelled by $\lambda = (\delta_a + \delta_{a-1}) \setminus \{1\}$, $\mathrm{Ind}_{W_L}^{W_1}(\lambda)$ contains all characters whose label is obtained by adding one box to the Young diagram of λ. Then $\mathrm{Ind}_{W_L}^W(\lambda) = \mathrm{Ind}_{W_1}^W \mathrm{Ind}_{W_L}^{W_1}(\lambda)$ can be computed as before. It ensues that the multiplicities of the characters labelled by $(\delta_a;\delta_{a-1})$ and by $(2a, 2a-5, \dots; 2a-3, 2a-7, \dots)$ are both 1. Again, induction gives the claim. The same reasoning applies for ${}^2D_n$ with n an odd square. □

2.3. **Cuspidal characters in quasi-isolated series.** We now study the action of automorphisms on cuspidal characters in quasi-isolated series of classical groups. The following result of Feit on Brauer trees will also provide some information on extendibility.

Lemma 2.3. *Let B be a block of a finite group H with cyclic defect. Let γ be an automorphism of H. If γ fixes some non-exceptional character in B, then it fixes every non-exceptional character in B.*

Proof. Assume that $\chi \in \mathrm{Irr}(B)$ is non-exceptional and fixed by γ. Then all nodes in the Brauer tree of B are fixed by γ by [8, Thm. 2.4], hence in particular all non-exceptional characters in B. □

Corollary 2.4. *Let $N \trianglelefteq H$ be finite groups with H/N solvable and of order prime to ℓ. Let χ, χ' be non-exceptional characters on the same ℓ-Brauer tree for H. Then χ extends to H if and only if χ' does.*

Proof. As H/N is solvable, there is a sequence of subgroups $N = U_1 \trianglelefteq \cdots \trianglelefteq U_r = H$ with U_i/U_{i-1} cyclic of prime order. Assume that χ extends to H, and let χ_i, $1 \leq i \leq r$, be compatible extensions of χ to U_i. Assume that we have extended χ' to a character χ_i' of U_i on the same ℓ-Brauer tree as χ_i. Since χ_i extends to H, it is invariant in U_{i+1}, hence by Lemma 2.3 the same is true for χ_i'. So χ_i' also extends to U_{i+1}, and clearly we may choose an extension χ_{i+1}' in the same ℓ-block as χ_{i+1}. So the claim follows by induction. □

Recall that Lusztig gives a partition $\mathrm{Irr}(G) = \coprod_s \mathcal{E}(G,s)$ of the set of irreducible characters of G into *rational Lusztig series* $\mathcal{E}(G,s)$ indexed by semisimple elements $s \in G^*$ up to conjugacy (see [7, Prop. 14.41]). Furthermore, as cuspidality is a uniform property,

by Jordan decomposition ([7, Thm. 13.25]) the cuspidal characters in the Lusztig series $\mathcal{E}(G,s)$ are in bijection with the cuspidal unipotent characters of $C_{G^*}(s)$ if $Z^\circ(C_{\mathbf{G}^*}(s))$ and $Z^\circ(\mathbf{G})$ have the same $\mathbb{F}_q$-rank, and there are none otherwise.

If $C^\circ_{\mathbf{G}^*}(s)^F$ is a product of classical groups, then it has a unique cuspidal unipotent character (see Table 1), so the cuspidal characters in $\mathcal{E}(G,s)$ form a single orbit under diagonal automorphisms. In particular, $\mathcal{E}(G,s)$ can contain cuspidal characters not fixed by some automorphism of G stabilising $\mathcal{E}(G,s)$ only if $C_{\mathbf{G}^*}(s)$ is not connected. Recall that s is *quasi-isolated in* $\mathbf{G}^*$ if $C_{\mathbf{G}^*}(s)$ is not contained in any proper F-stable Levi subgroup of $\mathbf{G}^*$.

Theorem 2.5. *Let G be quasi-simple of classical type B, C, D or ${}^2D$, let $s \in G^*$ be quasi-isolated and $\rho \in \mathcal{E}(G,s)$ cuspidal. Then there is a semisimple character $\chi \in \mathcal{E}(G,s)$ with the same stabiliser in* $\mathrm{Aut}(G)$ *as ρ.*

Proof. Let $\gamma \in \mathrm{Aut}(G)$. If $\mathcal{E}(G,s)$ is not stabilised by γ, then the claim holds trivially. So we may assume that $\mathcal{E}(G,s)$ is γ-stable. Moreover, if $C_{\mathbf{G}^*}(s)^F = C^\circ_{\mathbf{G}^*}(s)^F$ then $\mathcal{E}(G,s)$ contains a unique cuspidal and a unique semisimple character, and again we are done. Thus, as $|C_{\mathbf{G}^*}(s) : C^\circ_{\mathbf{G}^*}(s)|$ divides $|Z(\mathbf{G})|$, which is a 2-power and prime to p, we have in particular that q is odd. We discuss the remaining possibilities case-by-case.

The classes of quasi-isolated elements in $\mathbf{G}^*$ were classified by Bonnafé [1, Tab. 2]. The various rational types are worked out in Table 2. Our strategy of proof is as follows. In each case, $[C^\circ_{\mathbf{G}^*}(s), C^\circ_{\mathbf{G}^*}(s)]^F$ is a product of classical groups $G_1 \cdots G_r$. By the Jordan decomposition, the Lusztig series for s contains a cuspidal character if and only if each of these factors G_i has a cuspidal unipotent character ρ_i. If all factors are non-isomorphic, then for the factor of largest rank, say G_1, take a Zsigmondy prime ℓ as in Lemma 2.1. (Note that the exception $D_4(2)$ does not occur here as q is odd.) Then ρ_1 lies in an ℓ-block of cyclic defect, while all the other ρ_i are of ℓ-defect 0. In particular, the Jordan correspondent ρ of $\rho_1 \otimes \ldots \otimes \rho_r$ then also lies in a block of cyclic defect. As the Brauer trees in $\mathcal{E}_\ell(G,s)$ are just as for the Jordan corresponding unipotent blocks (see [9]), we conclude with Lemma 2.3 using the sequence of characters as in Lemma 2.1.

For example, in $G = \mathrm{Sp}_{2n}(q)$ centralisers of quasi-isolated elements in $G^* = \mathrm{SO}_{2n+1}(q)$ with $|C_{\mathbf{G}^*}(s) : C^\circ_{\mathbf{G}^*}(s)| = 2$ have types ${}^\delta D_d(q) B_{n-d}(q)$ for $1 \le d \le n$ and $\delta \in \{\pm\}$. These possess a cuspidal unipotent character $\rho_d \otimes \rho_{n-d}$ only if $d = a^2$ and $n - d = b(b+1)$ for some $a, b \ge 1$. Now for $a > b$ the cuspidal unipotent character ρ_d of ${}^\delta D_d(q)$ lies in a Brauer tree for primitive prime divisors ℓ of $q^{2a-1}+1$, while ρ_{n-d} is of ℓ-defect zero in $B_{n-d}(q)$; and for $a \le b$ we have that ρ_{n-d} lies in a Brauer tree for primitive prime divisors ℓ of $q^{2b}+1$, and ρ_d is of ℓ-defect zero.

Thus we are only left with those cases when $C_{G^*}(s)$ has two isomorphic quasi-simple factors, and these are the factors of largest rank. The corresponding lines are marked (1)–(5) in the last column of Table 2. We discuss them individually.

Cases 2, 4 and 5: Note that here cuspidal characters only arise when $\epsilon = -$, so when $q \equiv 3 \pmod 4$. But then q is not a square and so field automorphisms have odd order. In Case 2 the outer automorphism group of G is the direct product of the diagonal automorphism group A of order 2 with an odd order group of field automorphisms. So the latter must act trivially on all A-orbits. In Case 6 a Sylow 2-subgroup S of the outer

	2	${}^{\delta}D_d(q).D_{n-d}(q)$		2	$1 \le d < \frac{n}{2}$	$d=\square$, $n-d=\square$	
$(n \ge 4)$	2	${}^{\delta}D_{n/2}(q)^2$		2^2	n even	$n=2\square$	3)
	2	$D_{n/2}(q^2)$	$(2\times)$	2^2	n even	$n=2\square$	
	2	${}^{\delta}A_{n-1}(q).(q-\delta 1)$	$(2\times)$	2	n even	$\delta=-$, $n=\triangle$	
	4	${}^{\delta}D_d(q)^2.{}^{\epsilon}A_{n-2d-1}(q).(q-\epsilon 1)$		2^2	n even, $1 \le d < \frac{n}{2}$	$d=\square$, $\epsilon=-$, $n-2d=\triangle$	4)
	4	$D_d(q^2).{}^{-\epsilon}A_{n-2d-1}(q).(q+\epsilon 1)$	$(2\times)$	2^2	n even, $1 \le d < \frac{n}{2}$	$d=\square$, $\epsilon=+$, $n-2d=\triangle$	
	4	${}^{\delta}D_d(q)^2.A_{n-2d-1}(q).(q-1)$		4	n odd, $1 \le d < \frac{n}{2}$, $\epsilon=+$	never	
	4	${}^{2}D_d(q^2).{}^{2}A_{n-2d-1}(q).(q+1)$	$(2\times)$	4	n odd, $1 \le d < \frac{n}{2}$, $\epsilon=+$	$d=\square$, $n-2d=\triangle$	
	4	$D_d(q).{}^{2}D_d(q).{}^{2}A_{n-2d-1}(q).(q+1)$		2	n odd, $1 \le d < \frac{n}{2}$, $\epsilon=-$	$d=1$, $n-2=\triangle$	
	4	$D_d(q^2).A_{n-2d-1}(q).(q-1)$		2	n odd, $1 \le d < \frac{n}{2}$, $\epsilon=-$	never	
${}^{2}D_n(q)$	2	${}^{\delta}D_d(q).{}^{-\delta}D_{n-d}(q)$		2	$1 \le d < n/2$	$d=\square$, $n-d=\square$	
$(n \ge 4)$	2	$D_{n/2}(q).{}^{2}D_{n/2}(q)$		2	n even	never	
	2	${}^{2}D_{n/2}(q^2)$		2	n even	$n=2\square$	
	4	$D_d(q).{}^{2}D_d(q).{}^{\epsilon}A_{n-2d-1}(q).(q-\epsilon 1)$		2	n even, $1 \le d < \frac{n}{2}$	$d=1$, $\epsilon=-$, $n-2=\triangle$	
	4	${}^{2}D_d(q^2).{}^{-\epsilon}A_{n-2d-1}(q).(q+\epsilon 1)$		2	n even, $1 \le d < \frac{n}{2}$	$d=\square$, $\epsilon=+$, $n-2d=\triangle$	
	4	$D_d(q).{}^{2}D_d(q).A_{n-2d-1}(q).(q-1)$		2	n odd, $1 \le d < \frac{n}{2}$, $\epsilon=+$	never	
	4	$D_d(q^2).{}^{2}A_{n-2d-1}(q).(q+1)$		2	n odd, $1 \le d < \frac{n}{2}$, $\epsilon=+$	$d=\square$, $n-2d=\triangle$	
	4	${}^{\delta}D_d(q)^2.{}^{2}A_{n-2d-1}(q).(q+1)$		4	n odd, $1 \le d < \frac{n}{2}$, $\epsilon=-$	$d=\square$, $n-2d=\triangle$	5)
	4	${}^{2}D_d(q^2).A_{n-2d-1}(q).(q-1)$	$(2\times)$	4	n odd, $1 \le d < \frac{n}{2}$, $\epsilon=-$	never	

Here $\epsilon \in \{\pm\}$ is such that $q \equiv \epsilon 1$ (4), $\delta \in \{\pm\}$, ${}^{\pm}D_1(q)$ denotes a torus of order $q \mp 1$, $\square$ is a square, $\triangle$ a triangular number.

automorphism group of G consists of the diagonal automorphism group A_0 of order 4 extended by the graph-field automorphism of order 2. This is a dihedral group of order 8, as the graph automorphism of $\mathrm{Spin}^-_{2n}(q)$ centralises a subgroup $\mathrm{Spin}_{2n-1}(q)$ and so acts non-trivially on the centre. Clearly automorphisms of odd order must act trivially on any A_0-orbit they fix.

Now let $\gamma \in A$ be an automorphism of 2-power order moving a cuspidal character ρ. As there is just one class of elements s with the relevant centraliser, the image of ρ must lie in the A_0-orbit R of ρ, and A acts faithfully on R. The same then holds for the A-orbit of semisimple characters in $\mathcal{E}(G,s)$. In particular γ also moves some semisimple character. The same argument applies to Case 4 as the graph automorphism interchanges the two half-spin groups.

Case 1: Cuspidal characters occur if $n = 2m = 2a(a+1)$. Let $\mathbf{L} \leq \mathbf{G}$ be an F-stable Levi subgroup of twisted type ${}^2A_{n-1}(q).\Phi_2$ containing a Sylow 2-torus of $C_{\mathbf{G}^*}(s)$, with dual $\mathbf{L}^*$. Then $C_{\mathbf{L}^*}(s)$ is disconnected of type $C_{\mathbf{L}^*}(s)^F = ({}^2A_{m-1}(q).\Phi_2) \wr 2$. Let $\mathbf{G} \hookrightarrow \widetilde{\mathbf{G}}$ be our regular embedding, and $\widetilde{\mathbf{L}} = \mathbf{L}Z(\widetilde{\mathbf{G}})$. Let $\tilde{s}$ be an F-stable preimage of s in $\widetilde{\mathbf{G}}^*$. Then $C_{\widetilde{\mathbf{G}}^*}(\tilde{s}) = \tilde{\mathbf{H}}^2$ is connected with $\tilde{\mathbf{H}}$ of type $C_m(q)$, and $C_{\mathbf{L}^*}(\tilde{s}) = \widetilde{\mathbf{L}}_1^2$ is connected with $\widetilde{\mathbf{L}}_1^F = {}^2A_{m-1}(q).\Phi_2$. Jordan decomposition in $\tilde{G}$ maps $\mathcal{E}(\tilde{G},\tilde{s})$ to $\mathcal{E}(\tilde{H},1)\times\mathcal{E}(\tilde{H},1)$, and in $\tilde{L}$ it maps $\mathcal{E}(\tilde{L},\tilde{s})$ to $\mathcal{E}(\tilde{L}_1,1)\times\mathcal{E}(\tilde{L}_1,1)$. By Lemma 2.2(1) there is a unipotent character ψ_1 of $\tilde{L}_1$ such that $R^{\tilde{\mathbf{H}}}_{\widetilde{\mathbf{L}}_1}(\psi_1)$ contains the cuspidal unipotent character ρ_1 of $\tilde{H}$ and one other unipotent character χ_1 exactly once. As Lusztig induction of unipotent characters commutes with products, this implies that $R^{\tilde{\mathbf{H}}^2}_{\widetilde{\mathbf{L}}_1^2}(\psi_1^{\otimes 2})$ contains $\rho_1^{\otimes 2}$ and $\chi_1^{\otimes 2}$ exactly once.

As $\widetilde{\mathbf{L}}_1$ is of type A, all of its unipotent characters are uniform, and Jordan decomposition commutes with Deligne–Lusztig induction, so there is a unipotent character $\tilde{\psi}$ of $\widetilde{\mathbf{L}}$ such that $R^{\widetilde{\mathbf{G}}}_{\widetilde{\mathbf{L}}}(\tilde{\psi})$ contains the cuspidal character $\tilde{\rho} \in \mathcal{E}(\tilde{G},\tilde{s})$ and another unipotent character $\tilde{\chi}$ exactly once. The restriction of $\tilde{\psi}$ to L splits into two constituents ψ, ψ', and the restrictions of $\tilde{\rho}, \tilde{\chi}$ to G split into two constituents ρ, ρ' and χ, χ' respectively. As $R^{\mathbf{G}}_{\mathbf{L}}(\psi) + R^{\mathbf{G}}_{\mathbf{L}}(\psi')$ contains $\rho+\rho'$ and $\chi+\chi'$, we have that ρ, ρ' must occur with different multiplicities in $R^{\mathbf{G}}_{\mathbf{L}}(\psi)$, and similarly for χ, χ'. Now if γ is an automorphism of G not fixing ρ, then it cannot fix $R^{\mathbf{G}}_{\mathbf{L}}(\psi)$, hence neither χ. We conclude using the sequence of characters in Lemma 2.2.

Case 3: The argument here is very similar, using a Levi subgroup $\mathbf{L} \leq \mathbf{G}$ of type $D_2(q).{}^2A_{n-3}(q).\Phi_2$ with $C_{\mathbf{L}^*}(s)$ disconnected of type ${}^2A_{m-2}(q)^2.\Phi_2^4.2^2$ where $n = 2m = 2a^2$, and applying Lemma 2.2(2). □

Remark 2.6. By [5] the conclusion of Theorem 2.5 also holds for $\mathbf{G}$ of type A_n. For type C_n our result also follows from the recent work of Cabanes–Späth [6] and of Taylor [16].

3. Cuspidal characters in exceptional groups

We now turn to groups of exceptional type. Here, there exist Lusztig series containing more than one cuspidal character even for groups with connected centre, which makes the situation somewhat more involved.

3.1. **Trivial action.** We start by considering $\mathbf{G}$ simple simply connected of type E_6. Then $\mathbf{G}$ has a graph automorphism γ of order 2 which we may choose to commute with F.

Lemma 3.1. *Let $G = E_6(q)_{\mathrm{sc}}$ with $q \equiv 1 \pmod 3$, or $G = {}^2E_6(q)_{\mathrm{sc}}$ with $q \equiv -1 \pmod 3$ and $s \in G^*$ be semisimple. If $\mathcal{E}(G,s)$ is stabilised by the graph automorphism (resp. the graph-field automorphism) γ of G, then any automorphism commuting with γ acts trivially on non-trivial orbits of diagonal automorphisms in $\mathcal{E}(G,s)$.*

Proof. Let $X \subseteq \mathcal{E}(G,s)$ be a non-trivial orbit under the diagonal automorphisms, so $|X| = 3$. Write $q = p^f$ and first consider the case of ${}^2E_6(q)_{\mathrm{sc}}$. As $q \equiv -1 \pmod 3$ we have that q is not a square, so f is odd. Then $\mathrm{Out}(G) \cong \mathfrak{S}_3 \times C_f$, with the symmetric group $\mathfrak{S}_3$ inducing the diagonal automorphisms and the graph-field automorphism γ, and the cyclic group C_f inducing the field automorphisms. Now the $\mathfrak{S}_3$ factor necessarily acts faithfully on X. Thus any automorphism commuting with γ must centralise this $\mathfrak{S}_3$, so it acts trivially on X as claimed.

Now let $G = E_6(q)_{\mathrm{sc}}$. Then $\mathrm{Out}(G) \cong \mathfrak{S}_3 \times C_f$, with the first factor inducing diagonal and graph automorphisms, the second the field automorphisms if $p \equiv 1 \pmod 3$, respectively the product of the field automorphism by the graph automorphism if $p \equiv 2 \pmod 3$. We conclude as before. □

3.2. **Cuspidal characters in quasi-isolated series.** Next, we prove Theorem 1 for quasi-isolated series in exceptional type groups.

Theorem 3.2. *Let G be quasi-simple of exceptional type, $s \in G^*$ quasi-isolated and $\rho \in \mathcal{E}(G,s)$ cuspidal. Then there is a semisimple character $\chi \in \mathcal{E}(G,s)$ with the same stabiliser as ρ in $\mathrm{Aut}(G)$.*

Proof. The quasi-isolated elements were classified by Bonnafé [1]. We deal with the various possibilities case by case. First consider elements s with connected centraliser $\mathbf{C} = C_{\mathbf{G}^*}(s)$, for which we claim that cuspidal characters in $\mathcal{E}(G,s)$ are fixed by all automorphisms. If $s = 1$, so $\rho \in \mathcal{E}(G,1)$ is cuspidal unipotent then by results of Lusztig ρ is invariant under all automorphisms of G, see e.g. [11, Thm. 2.5], as is the semisimple character $1_G \in \mathcal{E}(G,1)$. Next assume that $s \neq 1$. If $\mathbf{C}$ has only components of classical type and $\mathbf{C}^{\circ F}$ does not involve ${}^3D_4$, then each such component has at most one cuspidal unipotent character and one semisimple character in any Lusztig series, and hence by the Jordan decomposition the same is true for $\mathcal{E}(G,s)$. The claim then follows trivially. Connected centralisers of quasi-isolated elements $s \neq 1$ with an exceptional component are of type E_6 in E_7, or of types $E_6.A_2$ or $E_7.A_1$ in type E_8. Now, the cuspidal unipotent characters of $E_6(q)$ lie on a Brauer tree for primes dividing Φ_9 together with the trivial character. Thus they lie on such a Brauer tree together with the semisimple character if they occur in quasi-isolated series of type E_6 or $E_6.A_2$. Our claim thus follows. Similarly two of the three cuspidal unipotent characters of ${}^2E_6(q)$ lie on the Φ_{18}-Brauer tree, and the third one is uniquely determined by its degree; and the two cuspidal unipotent characters of $E_7(q)$ lie on a Φ_{18}-Brauer tree together with the trivial character, and we conclude as before.

Thus we may now assume that $C_{\mathbf{G}^*}(s)$ is not connected. As centralisers of semisimple elements are connected in a group whose dual has connected centre, we have that $q \equiv 1 \pmod 3$ for $E_6(q)_{\mathrm{sc}}$, $q \equiv -1 \pmod 3$ for ${}^2E_6(q)_{\mathrm{sc}}$, and q is odd for $E_7(q)_{\mathrm{sc}}$, respectively. The various rational forms of the occurring types of disconnected centralisers $C_{\mathbf{G}^*}(s)$ with

cuspidal unipotent characters are collected in Table 3, depending on certain congruence conditions on q. The column labelled "$|\mathcal{E}(G,s)|$" gives the number of regular orbits (of length 3 or 2) under the group of diagonal automorphisms and of orbits of length 1 respectively, and similar the last column gives the same information for the subset of cuspidal characters. Note that we only need to concern ourselves with the regular orbits.

$\mathbf{G}^*$	$C_{\mathbf{G}^*}(s)^F$		q	$\lvert\mathcal{E}(G,s)\rvert$	$\lvert\mathcal{E}_{\mathrm{cusp}}(G,s)\rvert$
$E_6(q)$	${}^3D_4(q).\Phi_3.3$	$(2\times)$	$\equiv 1\ (3)$	8×3	2×3
${}^2E_6(q)$	${}^2A_2(q)^3.3$		$\equiv 2\ (3)$	$3\times 3+8\times 1$	1×3
	${}^2A_2(q^3).3$	$(2\times)$	$\equiv 2\ (3)$	3×3	1×3
	$D_4(q).\Phi_2^2.3$		$\equiv 2\ (3)$	$8\times 3+2\times 1$	1×3
	${}^3D_4(q).\Phi_6.3$	$(2\times)$	$\equiv 2\ (3)$	8×3	2×3
$E_7(q)$	${}^2E_6(q).\Phi_2.2$		$\equiv 1\ (2)$	30×2	3×2
	$D_4(q).A_1(q)^2.\Phi_2.2$		$\equiv 3\ (4)$	$20\times 2+18\times 1$	1×2
	${}^2A_2(q)^3.\Phi_2.2$		$\equiv 5\ (6)$	$9\times 2+9\times 1$	1×2

TABLE 3. Cuspidal characters in Lusztig series of quasi-isolated elements with disconnected centralisers in exceptional types

We consider these in turn, starting with $G=E_6(q)_{\mathrm{sc}}$. It can be checked by direct computation in Chevie (see [13]) that the two classes of quasi-isolated elements are invariant under the graph automorphism. In particular the corresponding Lusztig series must be invariant under the graph automorphism of G. Moreover, the $\tilde{G}$-orbits of cuspidal characters in those series $\mathcal{E}(G,s)$ are invariant by degree reasons. So our claim follows from Lemma 3.1. The situation and the arguments are entirely similar for the six quasi-isolated series in ${}^2E_6(q)_{\mathrm{sc}}$.

In $G=E_7(q)_{\mathrm{sc}}$, in the second case the cuspidal characters are contained in Brauer trees for Zsigmondy primes dividing Φ_6 (again note that q is odd, so in particular $q\neq 2$).

In the first case, let $\mathbf{L}^*\leq\mathbf{G}^*$ be the Levi subgroup of type $A_2(q).A_1(q^3).\Phi_3$ belonging to the nodes $1,2,3,5,7$ of the Dynkin diagram in the standard numbering. A Chevie-calculation shows that its dual Levi subgroup $\mathbf{L}\leq\mathbf{G}$ has disconnected centre, and $C_{\mathbf{L}^*}(s)=A_2(q).\Phi_2\Phi_3\Phi_6.2$. Application of [2, Thm. 3.2] gives that $R_{\tilde{\mathbf{L}}}^{\tilde{\mathbf{G}}}(\psi)$ contains the cuspidal character ρ with label ${}^2E_6[1]$ as well as the semisimple character in $\mathcal{E}(\tilde{G},\tilde{s})$ exactly once, where ψ is the semisimple character in $\mathcal{E}(\tilde{L},\tilde{s})$. So the claim holds for ρ by an argument as for Case 1 in the proof of Theorem 2.5. The two other cuspidal characters in this series lie on a Brauer tree for Zsigmondy primes dividing Φ_{18}.

Finally, for the last case, take a 6-split Levi subgroup $\mathbf{L}^*\leq\mathbf{G}^*$ for the nodes $2,5,7$, of type $A_1(q^3).\Phi_6^2$, whose dual again has disconnected centre, and with $C_{\mathbf{L}^*}(s)=(q^3+1)\Phi_6^2.2$. Then $R_{\tilde{\mathbf{L}}}^{\tilde{\mathbf{G}}}(\psi)$ contains the cuspidal character in $\mathcal{E}(\tilde{G},\tilde{s})$ as well as the semisimple character exactly once and we conclude as before. □

Corollary 3.3. *Let ρ be a cuspidal character of a quasi-simple group G of Lie type in a quasi-isolated series and assume we are neither in cases (3) or (4) of Table 2 nor in the first case of Table 3. Then some $\tilde{G}$-conjugate of ρ extends to its inertia group in the*

extension $\hat{G}$ of G by graph and field automorphisms. In particular ρ satisfies part (ii) of the inductive McKay condition from [12, Thm. 2.1].

Proof. By the proofs of Theorem 2.5 and 3.2 we have connected ρ via Brauer trees for primes $\ell > 3$ by or Deligne–Lusztig characters to a semisimple character χ in the same Lusztig series. According to [15, Prop. 3.4(c)] there exists a semisimple character χ' in the $\tilde{G}$-orbit of that satisfies the cited condition (ii) and thus in particular extends to its inertia group I in $\hat{G}$. Thus, the corresponding $\tilde{G}$-conjugate ρ' of ρ also has inertia group I. If ρ, χ (and hence ρ', χ') are connected by Brauer trees, ρ' extends to I by Corollary 2.4. In all other cases not excluded in the statement, all Sylow subgroups of $\hat{G}/G$ are cyclic, and so ρ' also extends. This is part (ii) of the inductive McKay condition. □

4. Proof of Theorem 1

We are now ready to prove our main result. Here, for finite groups $U \trianglelefteq V$ and $\chi \in \mathrm{Irr}(V)$, $\psi \in \mathrm{Irr}(U)$ we write $\mathrm{Irr}(U \mid \chi)$ for the constituents of $\chi|_U$, and $\mathrm{Irr}(V \mid \psi)$ for those of ψ^V.

Proof of Theorem 1. Let $\mathbf{G}$ be simple, simply connected and $F : \mathbf{G} \to \mathbf{G}$ such that $G = \mathbf{G}^F$ is quasi-simple. Let $\rho \in \mathrm{Irr}(G)$ be cuspidal and $s \in G^*$ such that $\rho \in \mathcal{E}(G, s)$. Let $\mathbf{L} \leq \mathbf{G}$ be a minimal F-stable Levi subgroup of $\mathbf{G}$ containing $C_{\mathbf{G}^*}(s)$. According to [16, Thm. 9.5] $R_{\mathbf{L}}^{\mathbf{G}}$ induces an $\mathrm{Aut}(G)_s$-equivariant bijection (Jordan decomposition) between $\mathcal{E}(G, s)$ and $\mathcal{E}(L, s)$, sending cuspidal characters to cuspidal characters and semisimple characters to semisimple ones. Here $\mathrm{Aut}(G)_s$ denotes the stabiliser in $\mathrm{Aut}(G)$ of $\mathcal{E}(G, s)$. Note that by construction s is quasi-isolated in $\mathbf{L}^*$. Thus we have reduced our question to the corresponding one for quasi-isolated series in L.

Let $\mathbf{L}_0 := [\mathbf{L}, \mathbf{L}]$. Then for $\chi \in \mathcal{E}(L, s)$ and $\chi_0 \in \mathrm{Irr}(L_0 \mid \chi)$ we have $\mathrm{Irr}(L \mid \chi_0) \cap \mathcal{E}(L, s) = \{\chi\}$. Thus χ is uniquely determined by χ_0 (given its Lusztig series). Moreover, χ is cuspidal, respectively semisimple if and only if χ_0 is, and L_0 is invariant under all automorphisms of L. Thus we have reduced our claim for the Lusztig series $\mathcal{E}(L, s)$ to the corresponding one for the Lusztig series $\mathcal{E}(L_0, s_0)$ of L_0, where $s_0 \in \mathbf{L}_0^*$ is the image of s under the natural epimorphism $\mathbf{L}^* \to \mathbf{L}_0^*$ induced by the embedding $\mathbf{L}_0 \leq \mathbf{L}$. Clearly s_0 is quasi-isolated in $\mathbf{L}_0$.

As $\mathbf{G}$ is simply connected, $\mathbf{L}_0$ is also of simply connected type, hence a direct product $\mathbf{L}_1 \cdots \mathbf{L}_r$ of F-orbits $\mathbf{L}_i$, $1 \leq i \leq r$, of simple components of $\mathbf{L}_0$. Correspondingly, we have $\mathbf{L}_0^F = L_1 \cdots L_r$ with quasi-simple finite groups of Lie type L_i. Any irreducible character χ of L_0 is an outer tensor product of irreducible characters χ_i of the L_i, which are cuspidal, respectively semisimple, if and only if χ is. Moreover, if $\chi \in \mathcal{E}(L_0, s_0)$ then $\chi_i \in \mathcal{E}(L_i, s_i)$, with s_i the image of s_0 under the epimorphism $\mathbf{L}_0^* \to \mathbf{L}_i^*$ induced by the embedding $\mathbf{L}_i \to \mathbf{L}_0$. Again, the s_i are quasi-isolated in $\mathbf{L}_i^*$. Thus we have reduced our statement to the case of cuspidal characters in quasi-isolated series of quasi-simple groups.

For groups of type A_n and ${}^2A_n$ the claim then follows from [5], for the other groups of classical type it is contained in Theorem 2.5 and for groups of exceptional type all relevant cases have been dealt with in Theorem 3.2. □

Example 4.1. Let $F' : \mathbf{G} \to \mathbf{G}$ be a Frobenius endomorphism commuting with F and such that $Z(G) \subset \mathbf{G}^{F'}$. Let $s \in G^*$ such that $\mathcal{E}(G, s)$ is stable under the field or graph-field automorphism σ of G induced by F'. Then any cuspidal character $\rho \in \mathcal{E}(G, s)$ is

σ-invariant. Indeed, by [3, Thm. 3.5] any such σ fixes the regular characters in $\mathcal{E}(G,s)$, hence the (Alvis–Curtis dual) semisimple characters, hence ρ by Theorem 1.

5. On the inductive McKay condition

We apply our previous result to verify the inductive McKay condition for several series of simple groups at suitable primes. Let still $\mathbf{G}$ be simple of simply connected type, with regular embedding $\mathbf{G} \hookrightarrow \widetilde{\mathbf{G}}$. By definition $\tilde{G}$ induces the full group of diagonal automorphisms of G. Let D be the group of automorphisms of $\tilde{G}$ induced by inner, graph and field automorphisms of G. We investigate the following property of characters $\tilde{\chi} \in \mathrm{Irr}(\tilde{G})$:

$$\textit{there exists } \chi \in \mathrm{Irr}(G \mid \tilde{\chi}) \textit{ which is } D_{\mathrm{Irr}(G|\tilde{\chi})}\textit{-stable;} \qquad (\dagger)$$

here $D_{\mathrm{Irr}(G|\tilde{\chi})}$ denotes the stabiliser in D of the set of irreducible characters of G below $\tilde{\chi}$. The following is known:

Lemma 5.1. *Regular and semisimple characters of $\tilde{G}$ satisfy* (†).

Proof. First let $\tilde{\chi} \in \mathrm{Irr}(\tilde{G})$ be a semisimple character. Then the claim is just [15, Prop. 3.4(c)]. Regular characters are the images of semisimple characters under the Alvis–Curtis duality (see [7, 14.39]), whose construction commutes with automorphism, so we conclude by the previous consideration. □

Proposition 5.2. *Let $G = {}^2E_6(q)_{\mathrm{sc}}$ for a prime power q and $\ell|(q+1)$ be a prime. Then any $\tilde{\chi} \in \mathrm{Irr}(\tilde{G} \mid \mathrm{Irr}_{\ell'}(G))$ satisfies* (†).

$C_{G^*}(s)$	q	$\lvert\mathcal{E}(G,s)\rvert$
${}^2A_2(q^3).3 \quad (2\times)$	$\equiv 2\ (3)$	3×3
$D_4(q).\Phi_2^2.3$	$\equiv 2\ (3)$	$8\times 3+2\times 1$
$A_1(q)^4.\Phi_2^2.3$	$\equiv 5\ (6)$	$4\times 3+4\times 1$
$A_1(q)^3.\Phi_2^3.3$	$\equiv 2\ (3)$	$2\times 3+2\times 1$
$A_1(q).\Phi_2^5.3$	$\equiv 2\ (3)$	2×3
$\Phi_2^6.3$	$\equiv 2\ (3)$	1×3

Table 4. Some ℓ'-series in ${}^2E_6(q)_{\mathrm{sc}}$, $\ell|(q+1)$

Proof. Let $\tilde{s} \in \tilde{G}^*$ be semisimple such that $\tilde{\chi} \in \mathcal{E}(\tilde{G}, \tilde{s})$, and let $s = \pi(\tilde{s})$. If $\tilde{\chi}$ restricts irreducibly to G, the claim holds trivially. So we may assume $C_{\mathbf{G}^*}(s)$ is not connected and hence that $3|(q+1)$. From the list of character degrees of ${}^2E_6(q)_{\mathrm{sc}}$ provided by Lübeck [10], we obtain the Table 4 of conjugacy classes of semisimple elements $s \in G^*$ with $C_{\mathbf{G}^*}(s)^F \neq C^\circ_{\mathbf{G}^*}(s)^F$ and such that $\mathcal{E}(G,s)$ contains characters of ℓ'-degree. As in Table 3 we also give the number of $\tilde{G}$-orbits in $\mathcal{E}(G,s)$.

If s is quasi-isolated, the claim follows with Lemma 3.1 as in the proof of Theorem 3.2. In the other three cases the non-invariant characters are either regular or semisimple, so they all satisfy (†) by Lemma 5.1. □

Thus we can prove our second main result:

Proof of Theorem 2. If $\ell|(q-1)$ the claim is contained in [12, Thm. 6.4(a)]. So now assume that $q \equiv -1 \pmod{\ell}$. For $S = {}^2E_6(q)$ all $\tilde{\chi} \in \mathrm{Irr}(\tilde{G} \mid \mathrm{Irr}_{\ell'}(G))$ satisfy property (†) by Proposition 5.2. As moreover here $D/\mathrm{Inn}(G)$ is cyclic, all such characters $\tilde{\chi}$ satisfy condition (ii)(1) in [12, Thm. 2.1] and thus the result follows from [12, Thm. 6.4(c)]. For the other families of groups, we have that $q \equiv -1 \pmod{\ell}$ with $\ell \equiv 3 \pmod 4$, so q is not a square. Thus S has no even order field automorphisms, whence $\mathrm{Out}(G)$ is cyclic. The claim then again follows from [12, Thm. 6.4(c)]. □

It was shown in [4, Cor. 7.3] that the inductive McKay condition holds for ${}^2E_6(q)$ at all primes $\ell \geq 5$ if $q \not\equiv -1 \pmod 3$, as well as for $E_7(q)$, $B_n(q)$ (and so $C_n(q)$) if q is even.

References

[1] C. Bonnafé, Quasi-isolated elements in reductive groups. *Comm. Algebra* **33** (2005), 2315–2337.

[2] M. Broué, G. Malle, J. Michel, Generic blocks of finite reductive groups. Astérisque **212** (1993), 7–92.

[3] O. Brunat, F. Himstedt, On equivariant bijections relative to the defining characteristic. *J. Algebra* **334** (2011), 150–174.

[4] M. Cabanes, B. Späth, Equivariance and extendibility in finite reductive groups with connected center. *Math. Z.* **275** (2013), 689–713.

[5] M. Cabanes, B. Späth, Equivariant character correspondences and inductive McKay condition for type A. *J. Reine Angew. Math.*, to appear, doi: 10.1515/crelle-2014-0104.

[6] M. Cabanes, B. Späth, Inductive McKay condition for finite simple groups of type C. arXiv:1612.03741, 2016.

[7] F. Digne, J. Michel, *Representations of Finite Groups of Lie Type*. Volume 21 of London Mathematical Society Student Texts. Cambridge University Press, Cambridge, 1991.

[8] W. Feit, Possible Brauer trees. *Illinois J. Math.* **28** (1984), 43–56.

[9] P. Fong, B. Srinivasan, Brauer trees in classical groups. *J. Algebra* **131** (1990), 179–225.

[10] F. Lübeck, Character degrees and their multiplicities for some groups of Lie type of rank < 9. Available at http://www.math.rwth-aachen.de/∼Frank.Luebeck/chev/DegMult/

[11] G. Malle, Extensions of unipotent characters and the inductive McKay condition. *J. Algebra* **320** (2008), 2963–2980.

[12] G. Malle, B. Späth, Characters of odd degree. *Ann. of Math. (2)* **184** (2016), 869–908.

[13] J. Michel, The development version of the CHEVIE package of GAP3. *J. Algebra* **435** (2015), 308–336.

[14] J. Olsson, Remarks on symbols, hooks and degrees of unipotent characters. *J. Combin. Theory Ser. A* **42** (1986), 223–238.

[15] B. Späth, Inductive McKay condition in defining characteristic. *Bull. Lond. Math. Soc.* **44** (2012), 426–438.

[16] J. Taylor, Generalised Gelfand–Graev representations in small characteristics. *Nagoya Math. J.* **224** (2016), 93–167.

FB Mathematik, TU Kaiserslautern, Postfach 3049, 67653 Kaiserslautern, Germany.
E-mail address: malle@mathematik.uni-kl.de